

# APPROPRIATE CARE MEASURES (RHQDAPU MEASURE SET)

RATES
<b>25 Measure RHQDAPU ACM</b>
<b>HF1: Discharge Instructions</b>
<b>HF2: Evaluation of LV Function</b>
<b>HF3: ACEI or ARB for LVSD</b>
<b>HF4: Smoking Cessation Counseling</b>
<b>HF ACM</b>
<b>AMI1: Aspirin at Arrival</b>
<b>AMI2: Aspirin at Discharge</b>
<b>AMI3: ACEI or ARB for LVSD</b>
<b>AMI4: Smoking Cessation Counseling</b>
<b>AMI5: Beta-blocker at Discharge</b>
<b>AMI7a: Fibrinolytic within 30 Minutes</b>
<b>AMI8a: PCI within 90 Minutes</b>
<b>AMI ACM</b>
<b>PN2: Pneumococcal Vaccination Given/Screened</b>
<b>PN3b: Initial Blood Culture Before First Antibiotic Dose (ED only)</b>
<b>PN4: Smoking Cessation Counseling</b>
<b>PN5c: First Antibiotic Dose within 6 Hours</b>
<b>PN6: Antibiotic Selection Consistent with Guidelines</b>
<b>PN7: Influenza vaccination given/screened</b>
<b>PN ACM</b>
<b>SCIP1: Abx within 1 Hr Before Incision or within 2 Hrs if Vancomycin or Quinolone is Used</b>
<b>SCIP2: Received Prophylactic Abx Consistent with Recommendations</b>
<b>SCIP3: Prophylactic Abx Discontinued within 24 Hrs of Surgery End Time or 48 Hrs for Cardiac Surgery</b>
<b>SCIP4: Controlled 6 AM Postoperative Serum Glucose - Cardiac Surgery</b>
<b>SCIP6: Appropriate Hair Removal</b>
<b>SCIP-CARD2: Pre-Admission Beta-blocker and Perioperative Period Beta-blocker</b>
<b>SCIP-VTE1: Recommended VTE Prophylaxis Ordered During the Admission</b>
<b>SCIP-VTE2: Received VTE Prophylaxis within 24 Hrs Prior to or After Surgery</b>
<b>SCIP ACM</b>

